

Negative order KdV equation with both solitons and kink wave solutions

Zhijun Qiao¹ and Jibin Li²

¹Department of Mathematics, The University of Texas-Pan American
1201 W University Drive, Edinburg, TX 78539, U.S.

²Department of Mathematics, Zhejiang Normal University, Jinhua, Zhejiang, 321004, China
Email: qiao@utpa.edu

Abstract

In this paper, we report an interesting integrable equation that has both solitons and kink solutions. The integrable equation we study is $(\frac{-u_{xx}}{u})_t = 2uu_x$, which actually comes from the negative KdV hierarchy and could be transformed to the Camassa-Holm equation through a gauge transform. The Lax pair of the equation is derived to guarantee its integrability, and furthermore the equation is shown to have classical solitons, periodic soliton and kink solutions.

PACS. 02.30.Ik C Integrable systems.

PACS. 05.45.Yv C Solitons.

PACS. 03.75.Lm C Tunneling, Josephson effect, Bose-Einstein condensates in periodic potentials, solitons, vortices, and topological excitations.

1 Introduction

Soliton theory and integrable systems play an important role in the study of nonlinear water wave equations. They have many significant applications in fluid mechanics, nonlinear optics, classical and quantum fields theories etc. Particularly in recent years, more focuses have been pulled to integrable systems with non-smooth solitons, such as peakons, cuspons, since the study of the remarkable Camassa-Holm (CH) equation with peakon solutions [2]. Henceforth, much progress have been made in the study of non-smooth solitons for integrable equations [4-20, 22, 27-33].

In this paper, we consider the following integrable equation

$$(\frac{-u_{xx}}{u})_t = 2uu_x, \quad (1)$$

which is actually the first member in the negative KdV hierarchy [29]. Equation (1) is proven equivalent to the Camassa-Holm (CH) equation: $m_t + m_x u + 2mu_x = 0, m = u - u_{xx}$ through a gauge transform (see Remark 1 in the paper). Therefore, we find a simpler reduced form of the CH equation. The Lax pair of the equation (1) is derived to guarantee its integrability, and furthermore the equation is shown to have classical solitons, periodic solitons and kink solutions.

2 Derivation of Equation (1) and Lax Representation

Let us consider the Schrödinger-KdV spectral problem:

$$L\psi \equiv \psi_{xx} + v\psi = \lambda\psi, \quad (2)$$

where λ is an eigenvalue, ψ is the eigenfunction corresponding to the eigenvalue λ , and v is a potential function. One can easily get the following Lenard operator relation:

$$K\nabla\lambda = \lambda J\nabla\lambda, \quad (3)$$

where $\nabla\lambda \equiv \frac{\delta\lambda}{\delta v} = \psi^2$ is the functional gradient of the spectral problem (2) with respect to v , $K = \frac{1}{4}\partial^3 + \frac{1}{2}(v\partial + \partial v)$ and $J = \partial$ are two Hamiltonian operators as known in the literature [1].

By setting $v = -\frac{u_{xx}}{u}$, we have the product form of operators K , $\mathcal{L}$, L and their inverses

$$\begin{aligned} K &= \frac{1}{4}u^{-2}\partial u^2\partial u^2\partial u^{-2}, \quad K^{-1} = 4u^2\partial^{-1}u^{-2}\partial^{-1}u^{-2}\partial^{-1}u^2, \\ \mathcal{L} &= \frac{1}{4}\partial^{-1}u^{-2}\partial u^2\partial u^2\partial u^{-2}, \quad \mathcal{L}^{-1} = 4u^2\partial^{-1}u^{-2}\partial^{-1}u^{-2}\partial^{-1}u^2\partial, \\ L &= \partial^2 + v = u^{-1}\partial u^2\partial u^{-1}, \quad L^{-1} = u\partial^{-1}u^{-2}\partial^{-1}u, \end{aligned}$$

where $\mathcal{L} = J^{-1}K$ and its inverse $\mathcal{L}^{-1} = K^{-1}J$ are the recursion operators for the positive order and negative order KdV hierarchy that we study below.

Now, according to the Lenard's operators K and J , we construct the entire KdV hierarchy, and then we show the integrability of the hierarchy through solving a key operator equation.

Let $G_0 \in \text{Ker } J = \{G \in C^\infty(\mathbb{R}) \mid JG = 0\}$ and $G_{-1} \in \text{Ker } K = \{G \in C^\infty(\mathbb{R}) \mid KG = 0\}$. We define the Lenard's sequence

$$G_j = \begin{cases} \mathcal{L}^j G_0, & j \in \mathbb{Z}, \\ \mathcal{L}^{j+1} G_{-1}, & j \in \mathbb{Z}, \end{cases} \quad (4)$$

where $\mathcal{L}, \mathcal{L}^{-1}$ are defined by Eq. (2). Therefore we generate a hierarchy of nonlinear evolution equations (NLEEs):

$$v_{t_k} = JG_k = KG_{k-1}, \quad \forall k \in \mathbb{Z}, \quad (5)$$

which is called the entire KdV hierarchy. We will see below that the positive order ($k \geq 0$) gives the regular KdV hierarchy usually mentioned in the literature [1], while the negative order ($k < 0$) produces some interesting equations gauge-equivalent to the Camassa-Holm equation [2]. Apparently, this hierarchy possesses the bi-Hamiltonian structure because of the Hamiltonian properties of K, J . Let us now give special equations in the entire KdV hierarchy (5).

- Choosing $G_0 = 2 \in \text{Ker } J$ (therefore $G_1 = u$) leads to the second positive member of the hierarchy (5):

$$v_{t_2} = \frac{1}{2}v_{xxx} + \frac{3}{2}vv_x, \quad (6)$$

which is exactly the well-known KdV equation. Here there is nothing new. Therefore, the positive order ($k \geq 0$) in the hierarchy (5) yields the regular KdV hierarchy usually studied in the literature [1].

- Now, let us find kernel elements $G_{-1} \in \text{Ker } K$ in order to get the negative member of the hierarchy (5). Due to the product form of K and K^{-1} , $G_{-1} = K^{-1}0$ has the following three seed solutions:

$$G_{-1}^1 = f(t_n)u^2, \quad G_{-1}^2 = g(t_n)u^2\partial^{-1}u^{-2}, \quad G_{-1}^3 = h(t_n)u^2\partial^{-1}u^{-2}\partial^{-1}u^{-2},$$

where $f(t_n), g(t_n), h(t_n)$ are three arbitrarily given functions with respect to the time variables t_n , but independent of x . They produce three iso-spectral ($\lambda_{t_k} = 0$) negative order KdV hierarchies of Eq. (5)

$$v_{t_k} = J\mathcal{L}^{k+1} \cdot G_{-1}^l, \quad l = 1, 2, 3, \quad k = -1, -2, \dots \quad (7)$$

When $k = -1$, their representative equations are:

$$\left(-\frac{u_{xx}}{u}\right)_{t_{-1}} = 2f(t_n)uu_x, \quad (8)$$

$$\left(-\frac{u_{xx}}{u}\right)_{t_{-1}} = g(t_n)(2uu_x\partial^{-1}u^{-2} + 1), \quad (9)$$

$$\left(-\frac{u_{xx}}{u}\right)_{t_{-1}} = h(t_n)(2uu_x\partial^{-1}u^{-2}\partial^{-1}u^{-2} + \partial^{-1}u^{-2}). \quad (10)$$

Remark 1. Apparently, the first one is differential and simpler and exactly recovers the equation (1) after setting $f(t_n) = 1$, which we focus on in the current paper. Actually, these three representative equations (8), (9), and (10) come from $v_{t_{-1}} = JG_{-1} = JK^{-1}0$ with $v = -\frac{u_{xx}}{u}$. Clearly, $v_{t_{-1}} = JK^{-1}0$ is equivalent to $KJ^{-1}v_t = 0$ ($t = t_{-1}$), that is,

$$\left(\frac{v_{txx}}{v_x}\right)_x + 4\left(\frac{vv_t}{v_x}\right)_x + 2v_t = 0. \quad (11)$$

This equation is exactly the one studied by Fuchssteiner [16] (see equations (7.1) and (7.22) there. Equations (7.1) in [16] has a typo and should be same as (11). From [16], the

Camassa-Holm (CH) equation is gauge-equivalent to equation (11) through some hodograph transformations (7.11) and (7.12) in [16]. In our paper, through using $v = -\frac{u_{xx}}{u}$ we further reduce equation (11) to a more simple form (i.e. equation (1)):

$$\left(-\frac{u_{xx}}{u}\right)_t = 2uu_x. \quad (12)$$

In other words, we found a very interesting fact that equation (12) can be viewed as a reduction form of the CH equation due to the above gauge-equivalence. In next section, we will solve this form.

In the paper [19], the author dealt with equation $(\partial^2 + 4v + 2v_x\partial^{-1})v_t = 0$ by using the positive KdV hierarchy approach, and all soliton solutions were given implicitly. This equation could be transformed to $(-u_{xx}/u)_t = 2uu_x$ through $v = -u_{xx}/u$, like we mentioned earlier in our paper, but, this is only one of three reductions. So, solutions of this equation can not give all solitons of our equation $(-u_{xx}/u)_t = 2uu_x$. In our paper, we present all solitons and kink solutions in a explicit form. Also, there is the connection of the first negative KdV equation $(\partial^2 + 4v + 2v_x\partial^{-1})v_t = 0$ with sine-Gordon [20]. But, the equation $(-u_{xx}/u)_t = 2uu_x$ we propose in the current paper is not equivalent to the sine-Gordon equation, because the sine-Gordon equation has only kink solution while our equation has both kink solutions and classical solitons.

Of course, we may generate higher order nonlinear equations by selecting different members in the hierarchy. In the following, we will see that all equations in the KdV hierarchy (5) are integrable. Particularly, the above three equations (8), (9), and (10) are integrable.

Let us return to the spectral problem (2). Apparently, the Gateaux derivative matrix $L_*(\xi)$ of the spectral operator L in the direction $\xi \in C^\infty(\mathbb{R})$ at point v is

$$L_*(\xi) \triangleq \left. \frac{d}{d\epsilon} \right|_{\epsilon=0} U(u + \epsilon\xi) = \xi \quad (13)$$

which is obviously an injective homomorphism, i.e. $U_*(\xi) = 0 \Leftrightarrow \xi = 0$.

For any given C^∞ -function G , one may consider the following operator equation [3] with respect to $V = V(G)$

$$[V, L] = L_*(KG) - L_*(JG)L. \quad (14)$$

Theorem 1. *For the spectral problem (2) and an arbitrary C^∞ -function G , the operator equation (14) has the following solution*

$$V = -\frac{1}{4}G_x + \frac{1}{2}G\partial, \quad (15)$$

where $\partial = \partial_x = \frac{\partial}{\partial x}$, and subscripts stand for the partial derivatives in x .

Proof: A direct substitution will complete the proof.

Theorem 2. Let $G_0 \in \text{Ker } J$, $G_{-1} \in \text{Ker } K$, and let each G_j be given through the Lenard sequence (4). Then,

1. each new vector field $X_k = JG_k$, $k \in \mathbb{Z}$ satisfies the following commutator representation

$$L_*(X_k) = [V_k, L], \quad \forall k \in \mathbb{Z}; \quad (16)$$

2. the entire KdV hierarchy (5), i.e.

$$v_{t_k} = X_k = JG_k, \quad \forall k \in \mathbb{Z}, \quad (17)$$

possesses the Lax representation

$$L_{t_k} = [V_k, L], \quad \forall k \in \mathbb{Z}, \quad (18)$$

where

$$V_k = \sum V(G_j) L^{(k-j-1)}, \quad \sum = \begin{cases} \sum_{j=0}^{k-1}, & k > 0, \\ 0, & k = 0, \\ -\sum_{j=k}^{-1}, & k < 0, \end{cases} \quad (19)$$

and $V(G_j)$ is given by Eq. (15) with $G = G_j$.

Proof: Let us only prove the case for $k < 0$. We have

$$\begin{aligned} [V_k, L] &= - \sum_{j=k}^{-1} [V(G_j), L] L^{k-j-1} \\ &= - \sum_{j=k}^{-1} (L_*(KG_j) - L_*(JG_j)L) L^{k-j-1} \\ &= - \sum_{j=k}^{-1} L_*(KG_j) L^{(k-j-1)} - L_*(KG_{j-1}) L^{k-j} \\ &= L_*(KG_{k-1}) - L_*(KG_{-1}) L^k \\ &= L_*(KG_{k-1}) = L_*(JG_k) \\ &= L_*(X_k). \end{aligned}$$

Noticing $L_{t_k} = L_*(v_{t_k})$, we have

$$L_{t_k} - [V_k, L] = L_*(v_{t_k} - X_k).$$

The injectiveness of L_* implies the second result holds.

So, the entire KdV hierarchy (5) has Lax pair and all equations in the hierarchy are therefore integrable. In particular, the KdV equation (6) has the Lax pair $L_{t_1} = [W_1, L]$ with $L = \partial^2 + v$ and $W_1 = \partial^3 + \frac{3}{2}v\partial + \frac{3}{4}v_x$, which was well-known in the literature [1].

Interesting thing is that the first member ($k = -1$) in the negative order KdV hierarchy (7) has the standard Lax representation $L_{t-1} = [V_{-1}^l, L]$ with $V_{-1}^l = (\frac{1}{4}G_{-1,x}^l - \frac{1}{2}G_{-1}^l\partial) L^{-1}$, $l = 1, 2, 3$, $L = \partial^2 + v = u^{-1}\partial u^2\partial u^{-1}$, and $L^{-1} = u\partial^{-1}u^{-2}\partial^{-1}u$. In particular, the negative KdV equation (8) possesses the following Lax form: $L_{t-1} = [V_{-1}^1, L]$ with

$$\begin{aligned} V_{-1}^1 &= \left(\frac{1}{4}G_{-1,x}^1 - \frac{1}{2}G_{-1}^1\partial \right) L^{-1} \\ &= \left(\frac{1}{2}uu_x - \frac{1}{2}u^2\partial \right) L^{-1} \\ &= -\frac{1}{2}u\partial^{-1}u. \end{aligned}$$

All of those negative members in the hierarchies (7) are integrable.

3 All traveling wave solutions of (1)

Let us now consider the traveling wave solution of equation (1) through a generic setting $u(x, t) = U(x - ct)$, where c is the wave speed. Let $\xi = x - ct$, then $u(x, t) = U(\xi)$. Substituting it into equation (1) yields

$$c\left(\frac{U''}{U}\right)' = 2UU'. \quad (20)$$

Integrating it once, we obtain the following standard cubic Hamiltonian system for $c \neq 0$:

$$\frac{dU}{d\xi} = y = \frac{\partial H}{\partial y}, \quad \frac{dy}{d\xi} = gU + \frac{1}{c}U^3 = -\frac{\partial H}{\partial U}, \quad (21)$$


where g is an integral constant, and the Hamiltonian function is

$$H(U, y) = \frac{1}{2}y^2 - \frac{1}{2}gU^2 - \frac{1}{4c}U^4. \quad (22)$$

When $gc \geq 0$, (21) has only one equilibrium point $O(0, 0)$. When $gc < 0$, (21) has three equilibrium points $O(0, 0)$ and $E_{1,2}(\pm\sqrt{|cg|}, 0)$. Write that

$$h_0 = H(0, 0) = 0, \quad h_1 = H(\pm\sqrt{|cg|}, 0) = \frac{1}{4}cg^2.$$

By qualitative analysis, we have the bifurcations of phase portraits of (21) in the (c, g) -parametric plan shown in Fig.1 (1-1)-(1-6).






Fig. 1 The change of phase portraits of (21) in the (c, g) -parameter plan.

(1-1) $g = 0, c > 0$. (1-2) $g > 0, c > 0$. (1-3) $g > 0, c < 0$. (1-4) $g = 0, c < 0$. (1-5) $g < 0, c < 0$. (1-6) $g < 0, c > 0$.

Next, we present the exact traveling wave solutions of (1) in an explicit form.

Case 1: $c > 0, g = 0$ (see Fig.1 (1-1)).

In this case, corresponding to the saddle point $O(0, 0)$ (21) reads as $y = \pm \frac{U^2}{\sqrt{2c}}$. Using the first equation of (21) and taking integration, we obtain

$$U(\xi) = \mp \frac{\sqrt{2c}}{\xi + \xi_0}, \quad \xi = x - ct, \quad (23)$$

where ξ_0 is an initial value of ξ . Clearly, when $\xi \rightarrow -\xi_0$, $U(\xi) \rightarrow \infty$. i.e., $U(\xi)$ is unbounded at $\xi = \xi_0$. Thus, we have two unbounded breaking wave solutions shown in Fig.2.




Fig. 2 The profiles of the functions (23) where $\xi_0 = -1$.

Case 2: $c > 0, g > 0$ (see Fig.1 (1-2)).

Corresponding to the saddle $O(0, 0)$ (21) reads as $y^2 = gU^2 + \frac{1}{2c}U^4$. Using the first equation of (21) to take integration, we obtain

$$U(\xi) = \pm \frac{8Acg}{A^2 e^{\sqrt{g}(\xi + \xi_0)} - 8cge^{-\sqrt{g}(\xi + \xi_0)}}, \quad (24)$$

where A is an integrant constant. When $A^2 = 8cg$, the functions defined by (24) are discontinuous at $\xi = -\xi_0$. The profiles of (24) like Fig.2.

Case 3: $g > 0, c < 0$ (see Fig.1 (1-3)).

There exist three equilibrium points of (21) at $E_{1,2}$ and $O(0, 0)$. O is a saddle point, $E_{1,2}$ are center points.

Corresponding to two homoclinic orbits, defined by $H(U, y) = 0$, we have the parametric representations

$$U(\xi) = \pm \sqrt{2|c|g} \operatorname{sech} \sqrt{\frac{|c|g}{2}} \xi. \quad (25)$$

They give two soliton solutions of (1).

For $h \in (\frac{1}{4}cg^2, 0)$, corresponding to two families of periodic orbits of (21), defined by $H(U, y) = h$, i.e., $y^2 = \frac{1}{2|c|}(4|c|h + 2g|c|U^2 - U^4) = \frac{1}{2|c|}(r_1^2 - U^2)(U^2 - r_2^2)$, where $r_1^2 = g|c| + \sqrt{g^2c^2 - 4|c|h}$, $r_2^2 = g|c| - \sqrt{g^2c^2 - 4|c|h}$, we obtain the parametric representations of periodic wave solutions of (1) as follows:

$$U(\xi) = \pm r_1 \operatorname{dn} \left(\frac{r_1}{\sqrt{2|c|}} \xi, \frac{\sqrt{r_1^2 - r_2^2}}{r_1} \right). \quad (26)$$

For $h \in (0, \infty)$, corresponding to the family of periodic orbits of (21), enclosing three equilibrium points defined by $H(U, y) = h$, we have the following parametric representation of periodic wave solutions of (1):

$$u(\xi) = r_1 \operatorname{cn} \left(\sqrt{\frac{(r_1^2 - r_2^2)}{2|c|}} \xi, \frac{r_1}{\sqrt{r_1^2 - r_2^2}} \right). \quad (27)$$

Case 4: $g \leq 0, c < 0$ (see Fig.1 (1-4), (1-5)).

In this case, the origin $O(0, 0)$ of (21) is an unique equilibrium point, which is a center. There exists a family of periodic orbits of (21), enclosing the origin. (1) has the same parametric representation of periodic wave solutions as (27).

Case 5: $g < 0, c > 0$ (see Fig.1 (1-6)).

In this case, there exist three equilibrium points of (21) at $E_{1,2}$ and $O(0, 0)$. O is a center, $E_{1,2}$ are saddle points. The heteroclinic orbits, defined by $H(u, y) = h_1$, have the parametric representations

$$U(\xi) = \pm \sqrt{c|g|} \tanh \left(\frac{\xi}{\sqrt{2|g|}} \right), \quad (28)$$

which gives a kink wave solution and an anti-kink wave solution of (1).

We see from (22) that $y^2 = \frac{1}{2c}(4ch + 2cgU^2 + U^4)$. For $h \in (0, h_1)$, it can be written as $y^2 = \frac{1}{2c}((z_1^2 - U^2)(z_2^2 - U^2))$, where $z_1^2 = |g|c + \sqrt{g^2c^2 - 4ch}$, $z_2^2 = |g|c - \sqrt{g^2c^2 - 4ch}$. Thus, the family of periodic orbits, defined by $H(u, y) = h$, has the parametric representation

$$u(\xi) = Z_2 \operatorname{sn} \left(\frac{z_1 \xi}{\sqrt{2c}}, \frac{z_2}{z_1} \right), \quad (29)$$

which gives a family of periodic wave solutions of (1).

4 Conclusions

In this paper, we reported an interesting property of integrable system: solitons and kink solutions can occur in the same integrable equation, and those solutions are given explicitly.

Within our knowledge, this is probably the first integrable example possessing such property. We found this equation in the negative order KdV hierarchy, which is gauge-equivalent to the CH equation. Since equation (1) has the Lax pair, we may try to get r -matrix structure of the constrained system of Lax equations, and parametric and algebro-geometric solutions [28], but that is beyond the scope of this paper. The symmetry of equations (8), (9), and (10) were already discussed in [23]. Recently, a two-fold integrable hierarchy associated with the KdV equation was given in [21]. About other negative order integrable hierarchies, such as the AKNS, the Kaup-Newell, the Harry-Dym, the Toda etc, one may see the literature [29].

Acknowledgment

The Qiao's work is partially supported by the U. S. Army Research Office under contract/grant number W911NF-08-1-0511 and by the Texas Norman Hackerman Advanced Research Program under Grant No. 003599-0001-2009.

References

- [1] M.J. Ablowitz and H. Segur, Solitons and Inverse Scattering Transform, SIAM, Philadelphia, 1981.
- [2] R. Camassa and D. D. Holm, An integrable shallow water equation with peaked solitons, Phys. Rev. Lett. 71(1993), 1661-1664.
- [3] C. W. Cao, Commutator representation of isospectral equations, Chin. Sci. Bull. 34(1989), 723-724.
- [4] A. Constantin and W. A. Strauss, Stability of peakons, Comm. Pure Appl. Math. 53(2000), 603-610.
- [5] A. Constantin and W. A. Strauss, Stability of the Camassa-Holm solitons, J. Nonlinear Sci. 12 (2002), 415C422.
- [6] A. Constantin, On the Cauchy problem for the periodic Camassa-Holm equation, J. Differential Equations, 141(1997), 218-235.
- [7] A. Constantin and J. Escher, Wave breaking for nonlinear nonlocal shallow water equations, Acta Math., 181(1998), 229C243.
- [8] A. Constantin and J. Escher, Well-posedness, global existence and blow-up phenomena for a periodic quasilinear hyperbolic equation, Comm. Pure Appl. Math., 51(1998), 475C504.

- [9] A. Constantin and D. Lannes, The hydrodynamical relevance of the Camassa-Holm and Degasperis-Procesi equations, *Arch. Ration. Mech. Anal.*, 192(2009), 165C186.
- [10] A. Constantin and H. P. McKean, A shallow water equation on the circle, *Comm. Pure Appl. Math.* 52(1999), 949C982.
- [11] A. Degasperis, M. Procesi, Asymptotic integrability, *Symmetry and perturbation theory* (Rome, 1998), World Sci. Publ., River Edge, NJ, 23 (1999).
- [12] H. R. Dullin, G.A. Gottwald and D.D. Holm, An integrable shallow water equation with linear and nonlinear dispersion, *Phys. Rev. Lett.* 87(2001), 194501-4504.
- [13] H. R. Dullin, G. A. Gottwald and D. D. Holm, Camassa-Holm, Korteweg-de-Vries-5 and other asymptotically equivalent equations for shallow water waves, *Fluid Dyn. Res.* 33(2003) 73-95.
- [14] H. R. Dullin, G. A. Gottwald and D. D. Holm, On asymptotically equivalent shallow water wave equations, *Physica D* 190(2004) 1-14.
- [15] A. Fokas and B. Fuchssteiner, Symplectic structures, their Baecklund transformation and hereditary symmetries, *Physica D* 4(1981), 47-66.
- [16] B. Fuchssteiner, Some tricks from the symmetry-toolbox for nonlinear equations: generalizations of the Camassa-Holm equation, *Phys. D* 95(1996), 229-243.
- [17] D. D. Holm and R. I. Ivanov, Two-component CH system: Inverse Scattering, Peakons and Geometry, *arXiv:1009.5374*.
- [18] D. D. Holm and R. I. Ivanov, Multi-component generalizations of the CH equation: Geometrical Aspects, Peakons and Numerical Examples, *arXiv:1009.5368*.
- [19] A. N. W. Hone, The associated Camassa-Holm equation and the KdV equation, *J. Phys. A: Math. Gen.* 32(1999), L307-L314.
- [20] A. N. W. Hone and J.P.Wang, Prolongation algebras and Hamiltonian operators for peakon equations, *Inverse Problems* 19(2003), 129-145.
- [21] A. Kundu, Exact accelerating soliton in nonholonomic deformation of the KdV equation with two-fold integrable hierarchy, *J. Phys. A: Math. Theor.* 41(2008) 495201.
- [22] R. S. Johnson, Camassa-Holm, Korteweg-de Vries and related models for water waves, *J. Fluid Mech.* 455(2002), 63-82.
- [23] S. Y. Lou, Symmetries of the KdV equation and four hierarchies of the integrodifferential KdV equation, *J. Math. Phys.* 35(1994) 2390-2396.

- [24] H. P. McKean, The Liouville correspondence between the Korteweg-de Vries and the Camassa-Holm hierarchies, *Comm. Pure Appl. Math.* 56(2003), 998-1015.
- [25] Z. J. Qiao, A new integrable equation with cuspons and W/M-shape-peaks solitons, *J. Math. Phys.* 47(2006), 112701-9.
- [26] Z. J. Qiao, New integrable hierarchy, cuspons, one-peak solitons, and M/W-shape-peak solutions, *Journal of Mathematical Physics* 48(2007), 082701-20.
- [27] Z. J. Qiao, The Camassa-Holm hierarchy, N-dimensional integrable systems, and algebro-geometric solution on a symplectic submanifold, *Communications in Mathematical Physics* 239(2003), 309-341.
- [28] Z. J. Qiao, Generalized r -matrix structure and algebro-geometric solutions for integrable systems, *Rev. Math. Phys.* 13(2001), 545-586.
- [29] Z. J. Qiao, Finite-dimensional Integrable System and Nonlinear Evolution Equations, (English), 230 pages, ISBN 7-04-010516-0, Chinese National Higher Education Press, Beijing, PR China, 2002.
- [30] Z. J. Qiao and S. Li, A new integrable hierarchy, parametric solution, and traveling wave solution, *Mathematical Physics, Analysis and Geometry* 7(2004), 289-308.
- [31] Z. J. Qiao and G. Zhang, On peaked and smooth solitons for the Camassa-Holm equation, *Europhys. Lett.* 73(2006), 657-663.
- [32] Z. J. Qiao and W. Strampp, Negative order MKdV hierarchy and a new integrable Neumann-like system, *Physica A* 313(2002), 365-380.
- [33] V. Vakhnenko and E. Parkes, Periodic and solitary-wave solutions of the Degasperis-Procesi equation, *Chaos Solitons Fractals* 20(2004), 1059-1073.